



Outcomes of complex mitral valve repair through right mini-thoracotomy using artificial chordae

Vinh Duc An Bui¹, Dinh Hoang Nguyen^{2,*}, Chuong Tran Viet Pham², Thuan Quang Phan², Thanh Van Thai Nguyen³, Ngoc-Minh Vuong³, Thao Nhat Le², Thang Duc Ho², Tam Nhat Minh Ung⁴, Dung Hung Van⁵, and Nam Hoai Nguyen³

¹Department of Thoracic and Cardiovascular Surgery, Hue Central Hospital, Hue, Vietnam

²Department of Adult Cardiovascular Surgery, University Medical Center Ho Chi Minh City, Ho Chi Minh City, Vietnam

³Department of Cardiothoracic and Vascular Surgery, Faculty of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Vietnam

⁴Faculty of Medicine, University of Debrecen, Debrecen, Hungary

⁵Heart Institution of Ho Chi Minh City, Ho Chi Minh City, Vietnam

Abstract

Introduction: Mitral regurgitation (MR) affects 2%–3% of the global population. The rising incidence of degenerative causes in developing countries leads to an increase in the number of surgeries for degenerative MR. In this study, we aimed to evaluate the safety, feasibility, and effectiveness of minimally invasive mitral repair (MIMR) using artificial chordae in complex cases of mitral valve regurgitation.

Method: This is a retrospective analysis of MIMR through right mini-thoracotomy approach with intermediate-to-complex level of repair difficulty. We analyzed the demographic information, echocardiography results, surgical details, and mid-term survival outcomes.

Results: 49 patients underwent surgery with a mean age of 49.1 ± 13.3 years. 34 patients (69.39%) were classified as having intermediate level of MR complexity, four patients (8.2%) had a complex mitral valve score, and 11 patients (22.4%) were diagnosed with Barlow's disease. We performed a ring annuloplasty in all patients, with additional repair techniques including edge-to-edge (10.2%), commisuroplasty (26.5%), and interscallop indentation closure (22.5%). There were no in-hospital deaths. We recorded three cases with minor complications: one case of pleural effusion, one case of low cardiac output syndrome, and one case of acute lower limb compartment syndrome. Cumulative recurrent MR rates measured at 12-mon, 24-mon and 36-mon are 92%, 88%, and 84%, respectively.

Conclusions: MIMR of complex mitral valves using artificial chordae was effective, with low complication and recurrent rates. Optimal results can be achieved through proper lesion evaluation and choosing the appropriate repair strategies using artificial chordae combining with edge-to-edge, commisuroplasty or indentation closure techniques.

Keywords: mitral valve prolapse; mitral valve insufficiency; artificial chordae; complex mitral valve repair; minimally invasive surgical procedures

Received: Jun 22, 2023 / Revised: Jul 12, 2023 / Accepted: Aug 18, 2023

*Corresponding author: Dinh Hoang Nguyen. Department of Adult Cardiovascular Surgery, University Medical Center Ho Chi Minh City, Ho Chi Minh City, Vietnam. E-mail: dinh.nh@umc.edu.vn

Copyright © 2024 MedPharmRes. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Mitral regurgitation (MR) affects 2%–3% of the global population [1]. The incidence of MR caused by degenerative factors is rising in developing countries, while rheumatic valve disease remains the primary cause of MR [1],[2]. One study in Vietnam shows there is a decrease in rheumatic mitral valve surgery from 95% to 81%, and an increase in the number of surgeries due to degenerative causes from 1.6% to 16%, and in the rate of surgery for degenerative MR from 11.4% to 38.5% [3].

The degenerative mitral valve pathomechanism affects the posterior leaflet or annulus and can typically be repaired through conventional repair techniques. Posterior leaflet prolapse can be corrected through leaflet resection or artificial chordae implantation while correcting the anterior or bi-leaflet prolapse requires more complex procedures and strategies. We aim to examine the safety and feasibility of minimally invasive mitral repair using artificial chordae in intermediate-to-complex valve regurgitation. Our goal is to determine whether such operations can be effectively applied, despite the challenges posed by complex valve pathologies.

2. MATERIALS AND METHODS

A retrospective analysis was performed on patients who underwent minimally invasive mitral valve repair using artificial chordae via right mini-thoracotomy at one center in Vietnam between April 2016 and April 2022. The indication for mitral valve surgery followed the current American College of Cardiology/American Heart Association (ACC/AHA) guidelines for the management of valvular heart disease [4]. A single, primary, cardiovascular surgeon, who had overcome the learning curve of mitral valve repair [5], and supported by his assistants, performed all the operations. The selection of the patient population was based on the Anyanwu complexity score. Intermediate and complex repair cases were selected. Furthermore, demographic information, echocardiographic findings, operative characteristics, surgical outcomes, and mid-term survival rates of the patients were

collected and analyzed.

The Anyanwu complexity score [6] was used to categorize the valve repair into three levels of complexity: simple (score \geq 1), intermediate (score \geq 2–4), and complex (score \geq 5). Residual MR was defined as the MR of more than moderate severity at the time of transferring the patient from the operating room. Recurrent MR was defined as the occurrence of MR after an initial absence of the condition at the time of discharge from the operating room.

The data was analyzed using Microsoft Excel 2016 and SPSS version 16.0. Descriptive analysis was used to analyze the patient's demographic and clinical characteristics and presented as mean \pm SD for continuous variables and frequency and percentage for categorical variables. Survival, freedom from reoperation, and recurrent MR were evaluated using Kaplan-Meier analysis.

3. RESULTS

49 patients with intermediate-to-complex degenerative MR underwent minimally invasive mitral valve repair using artificial chordae via right mini-thoracotomy. The mean age was 49.1 \pm 13.3 years, with a male-to-female ratio of 3.1:1. The mean EuroSCORE II was 1.12 \pm 0.99%. All cases of MR in the study were classified as severe and categorized as Carpentier's functional classification type II, with no cases of moderate-to-severe tricuspid regurgitation requiring repair observed (Tables 1–4, Figs. 1 and 2).

4. DISCUSSION

The current guidelines on mitral valve surgery emphasize repairing the valve instead of replacing it when possible [4]. The feasibility of repair, which is highly affected by the intricacy of the lesion and the surgeon's expertise, plays a role in determining if surgery is needed.

To ensure a successful MR repair and assess its complexity, it's crucial to use a proper grading method to assist in choosing appropriate techniques. Consultation with experts is encouraged, if needed. Surgeon experience is commonly assessed based on his years in practice. Li et al. [7] found

Table 1. Characteristics of previous pregnancy by study group

Characteristic	Value (N=49)
Demographic characteristics	
BMI (kg/m ²)	22.6±2.7
BSA (m ²)	1.65±0.17
NYHA I-II	44 (89.8%)
Atrial fibrillation	4 (8.2%)
Hypertension	22 (44.9%)
Diabetes mellitus	6 (12.2%)
Echocardiography characteristics	
Vena contracta (mm)	8.4±1.8
Annular mitral diameter (mm)	40.2±4.5
LVEDD (mm)	59.9±6.0
LVESD (mm)	37.0±5.2
LA diameter (mm)	47.3±8.5
LA volume (mL)	144.8±57.7
LVEF (%)	63.4±8.8
TAPSE (mm)	25.8±4.2
sPAP (mmHg)	38.9±20.4
Valvular prolapse at	
A ₁	6 (12.2%)
A ₂	22 (44.9%)
A ₃	15 (30.6%)
P ₁	16 (32.7%)
P ₂	23 (46.9%)
P ₃	12 (24.5%)
Complexity score ¹⁾	
Intermediate	45 (91.8%)
Complex	4 (8.2%)
Etiology of mitral valve	
FED	38 (77.6%)
Barlow	11 (22.4%)

¹⁾ Based on Anyanwu's score. BMI, body mass index; BSA, body surface area; NYHA, New York Heart Association; LVEDD, left ventricular end-diastolic diameter; LVESD, left ventricular end-systolic diameter; LA, left atrial; LVEF, left ventricular ejection fraction; TAPSE, tricuspid annular plane systolic excursion; sPAP, systolic pulmonary artery pressure; A, anterior; P, posterior; FED, fibroelastic deficiency.

that it generally takes between 50 to 200 operations for surgeons to surpass the learning curve associated with mitral valve repair rates, with some variation among individual surgeons. In our previous publication, when the primary surgeon has successfully surpassed the lower threshold of 90 operations, he has overcome the learning curve for mitral repair rates, thus enabling him to consider repairing more complex mitral valve lesions [5]. In general, the complexity of a mitral valve repair is considered simple if only the posterior

Table 2. Surgical characteristics

Surgical characteristic	Value (N=49 [%])
Artificial chordae at	
A ₁	2 (4.1)
A ₂	20 (40.8)
A ₃	9 (18.4)
P ₁	2 (4.1)
P ₂	17 (34.7)
P ₃	9 (18.4)
Ring annuloplasty	
28 mm	11 (22.5)
30 mm	14 (28.6)
32 mm	17 (34.7)
34 mm	6 (12.2)
36 mm	1 (2.0)
Additional repair technique	
Edge-to-edge technique	5 (10.2)
Commisuroplasty	13 (26.5)
Interscallop indentation closure	11 (22.5)
Concomitant procedures	
Atrial fibrillation ablation	4 (8.2)
ASD closure	1 (2.0)

The mean aortic clamp time was 92.6±20.6 min. The mean CPB time was 201.1±67.7 min. A, anterior; ASD, atrial septal defect; P, posterior; CPB, cardiopulmonary bypass.

Table 3. Postoperative features

Postoperative feature	Value (N=49, [%])
In-hospital mortality	0 (0.0)
Prolonged ventilation time	6 (12.2)
Low-cardiac output syndrome	1 (2.0)
Re-exploration	1 (2.0)
Temporary renal replacement therapy	0 (0.0)
Limb compartment syndrome	1 (2.0)
New onset of atrial fibrillation	9 (18.4)
Mitral re-regurgitation	
None/trivial	40 (81.6)
Moderate	9 (18.4)
Severe	0 (0.0)

Table 4. Mid-term outcomes

Mid-term outcome	Value (N=49 [n])
Overall mortality	1 (2.0)
Mitral reoperation	0 (0.0)
Follow-up mitral re-regurgitation	
None/trivial	43 (87.8)
Moderate to severe	6 (12.2)

The follow-up period was 27.8±16.9 mon.




Fig. 1. The Kaplan-Meier survival curve showed that the survival estimates were 100% at 12 and 24 mon and 96% (89% to 100%) at 36 mon.




Fig. 2. The freedom from recurrence MR after 12, 24, and 36 mon was 92% (84% to 100%), 88% (79% to 99%), and 84% (72% to 98%), respectively.

leaflet is prolapsed; and is complex if the anterior leaflet or both leaflets are prolapsed [8]. Different authors have evaluated the complexity of mitral valve repair through various grading scores [9],[10]. However, Anyanwu's system, based on echocardiography features, the location of leaflet prolapse, valve tissue calcification, and limited leaflet mobility is widely considered practical and comprehensive [6]. In our study, we used the complexity score of Anyanwu, as it has also been utilized by Nakayama [11].

Eleven patients (22.4%) were diagnosed with Barlow's disease, a complex condition characterized by excessive myxomatous tissue in the leaflets, bi-leaflet prolapse, and severe annular dilation, which posed significant challenges. Four patients (8.2%) presented with complex valvular lesions based on Anyanwu's grade. All these complex forms of mitral valves were repaired using a combination of ring annuloplasty and other repair techniques in which artificial chordae were used as the principal technique. Another important aspect of Barlow valve repair is mitral annular disjunction, a condition characterized by a separation between the atrial-mitral valve junction and ventricular attachment [4]. This can be observed during echocardiography video perioperatively. In these cases, the mitral annulus may have a flatter and more elliptical shape, with reduced height and loss of the saddle shape. Therefore, when implanting an annular mitral ring, using a complete rigid ring with a saddle shape is crucial to provide a more favorable and uniform distribution of forces, reducing strain on the mitral leaflet and improving leaflet coaptation geometry [12]. According to Rankin et al. [13], almost all patients with mitral disease can be effectively treated using artificial chord repair, even in complex cases. Our results showed that none of the patients had residual or recurrent MR at discharge. Nine patients (18.4%) had moderate MR according to postoperative echocardiography, and there was no conversion to valve replacement.

In the past, the technique for repairing posterior leaflet prolapse involved making a triangular or quadrangular resection on the posterior leaflet, followed by leaflet plication. Sliding-plasty techniques can reduce the risk of Systolic Anterior Motion (SAM) while ensuring the repair is effective [14], but may result in decreased mobility of the posterior leaflet.

To overcome this issue, there has been a trend towards using artificial chordae for repairing extensive or multiple segments of prolapse of the posterior leaflet, especially in the P2 scallop. This procedure is considered more straightforward and allows for adjusting the height of the posterior leaflet, reducing the risk of SAM [15].

For repairing anterior leaflet prolapse or bi-leaflet prolapse, there is a general consensus among surgeons to use artificial chordae for all anterior prolapses. We would like to emphasize that bi-leaflet prolapse is a complex condition to repair. Castillo et al. [16] proposed a two-step process: first, repairing the posterior leaflet, followed by inserting a ring annuloplasty, and then conducting a saline test to evaluate the regurgitation state. Most injuries, including complex lesions, can be thoroughly evaluated at this stage, and various valve repair techniques, such as artificial chordae, commissurotomy, and edge-to-edge stitch, could be applied. According to Koprivanac et al. [17], in bi-leaflet prolapse, an annuloplasty ring is appropriate for repair if the prolapse is large and even, and neo-chordae implantation is suitable if the prolapse is asymmetrical.

In five cases (10.2%), we employed the Edge-to-Edge technique as a backup solution in case of an inadequate repair. This method is favored due to its transparency and efficiency, especially in challenging situations. The surgical procedure involved stitching the loose edges of the leaflets at the site of regurgitation, resulting in a valve with two orifices when the regurgitation originates from the middle folds, as described by Alfieri et al. [18].

We employed commissuroplasty in 13 cases (26.5%). Although less common than anterior and posterior leaflet prolapse, commissural prolapse is not uncommon and often occurs as residual lesions following the primary repair. Diagnosis can be challenging, as it may be missed in one-third of cases and difficult to detect through preoperative echocardiography [19]. The typical repair method is commissural closure, which does not result in significant flow restriction [19]. In addition, 11 patients (22.5%) underwent interscallop indentation closures. Indentation can develop as the leaflet volume expands, which can be easily identified through a saline test and can be easily repaired with a single stitch.

Our surgical results showed a mean aortic clamp time of 92.6 ± 20.6 min and a mean cardiopulmonary bypass (CPB) time of 201.1 ± 67.7 min, which were longer than the results achieved by Nakayama [11]. The author repaired 141 cases of MR using a mini-invasive approach via right thoracotomy (81% classified as intermediate to complex), and outcomes showed a CPB time of 144 ± 36 min and a cross-clamp time of 113 ± 35 min [11].

There were no in-hospital deaths in our study. During the postoperative period, one case of pleural effusion required re-exploration via the same thoracic incision. Another case of postoperative low cardiac output syndrome was recognized, caused by low blood pressure and a decline in left ventricular function. An intra-aortic balloon pump was inserted and successfully removed. The patient was then discharged from the hospital successfully. Nakayama did not encounter any cases of low cardiac syndrome in his record [11].

A case of postoperative lower limb compartment syndrome was recorded in a patient who underwent peripheral CPB with a 22-Fr femoral artery cannula, leading to an extended CPB duration of 216 min. The syndrome was diagnosed 10 h post-surgery and presented symptoms of severe pain, coldness, tenderness, paresthesia, and weak peripheral pulses. The limb was saved with a fasciotomy. The patient's high body mass index, large CPB cannula size, extended operation time, and Kawashima's Type D vasculature [20] were identified as risk factors. The incidence of limb compartment syndrome was extremely low. Out of 2,645 patients who underwent minimally invasive cardiac surgery, only two cases of lower extremity compartment syndrome (0.08%) were documented [21]. The rareness and uniqueness of this patient's situation have been reported in our previous case report publication [22].

In this study, freedom from recurrence of MR was 92% at 12-mon, 88% at 24-mon, and 84% at 36-mon. These results suggest a low incidence of recurrent MR but inferior to those found in the outcomes of Castillo et al. [16], which involved 188 patients with degenerative mitral valve disease and either isolated anterior leaflet or bi-leaflet prolapse. Research by David et al. [23] indicates that early recurrent MR is often caused by technical errors or insufficient repair, while late

recurrent MR is mainly due to degenerative progression. During follow-up, no reoperations due to MR recurrence were found in our study. According to Chitwood [24], mitral valve repairs can fail, despite being performed by skilled surgeons. The reasons behind these failures can be attributed to the progression of the disease, endocarditis, and technical issues such as the dehiscence of the prosthesis and the tearing of sutures.

Our results revealed that the survival rate measuring at 24-mon and 36-mon was 100% and 96%, respectively. Only one death was recorded during the follow-up. These results are considered favorable when compared to previous studies. For example, Nakayama reported 1-year and 3-year survival rates of 99.2% and 98.1%, respectively [11]. Castillo and his colleagues [16] found a cumulative survival rate of 96.3% at 4-year in patients in case of bi-leaflet prolapse. The author's predominant repair technique was artificial chordae or loop in 49% of operations. Overall, these findings suggest that our results demonstrate a satisfactory level of patient survival compared to previous studies.

The current study has several limitations. Firstly, its retrospective design may introduce biases and limitations in data collection. Additionally, the relatively small sample size and single-center setting raise concerns about the generalizability of the findings. Another limitation is that the Anyanwu complexity classification used in the study may not be comprehensive enough as it did not consider certain factors such as the presence of mitral annular disjunction. Moreover, since the study was conducted by a single surgeon, there may be biases due to differences in preferred repair strategies compared to other surgeons. To address these limitations, future research should involve larger prospective studies conducted in multiple centers with different surgeons, using different complexity score systems to enhance the reliability of the evidence.

5. CONCLUSION

In conclusion, our study demonstrates that minimally invasive repair of complex mitral valves using artificial chordae is a feasible approach with low morbidity rates and recur-

rence of MR. The success of this repair technique depends on the accurate evaluation of the lesion's complexity using proper grading system and perioperative valve analysis. By combining different repair techniques with the correct repair strategies, it could enhance the success rate of MR repair significantly.

Acknowledgements

The authors would like to thank the Faculty of Pharmacy, University of Medicine and Pharmacy at Ho Chi Minh City, who supported and facilitated the authors to complete this study.

Funding sources

Vinh D.A. Bui was funded by Vingroup JSC and supported by the Master, Ph.D. Scholarship Programme of Vingroup Innovation Foundation (VINIF), Institute of Big Data (code VINIF.2022.TS146).

Conflict of interest

No potential conflict of interest relevant to this article was reported.

ORCID

Vinh Duc An Bui

<https://orcid.org/0000-0002-3833-5709>

Dinh Hoang Nguyen

<https://orcid.org/0000-0002-6769-0849>

Chuong Tran Viet Pham

<https://orcid.org/0000-0002-8262-7848>

Thuan Quang Phan

<https://orcid.org/0000-0003-4523-7442>

Thanh Van Thai Nguyen

<https://orcid.org/0000-0003-1383-8606>

Ngoc-Minh Vuong

<https://orcid.org/0000-0002-2199-2569>

Thao Nhat Le

<https://orcid.org/0000-0002-2994-9776>

Thang Duc Ho

<https://orcid.org/0000-0002-7467-3975>

Tam Nhat Minh Ung

<https://orcid.org/0000-0002-7126-9109>

Dung Hung Van

<https://orcid.org/0000-0002-9652-1825>

Nam Hoai Nguyen

<https://orcid.org/0009-0009-5545-090X>

Authors' contributions

Conceptualization: DH Nguyen, TQ Phan.

Data curation: VDA Bui, TN Le, TD Ho.

Formal analysis: VDA Bui, TVT Nguyen.

Methodology: DH Nguyen, NH Nguyen.

Software: VDA Bui, CTV Pham.

Validation: DH Nguyen, CTV Pham.

Investigation: DH Van, NH Nguyen.

Writing - original draft: VDA Bui.

Writing - review & editing: VDA Bui, DH Nguyen, CTV Pham, TQ Phan, TVT Nguyen, NM Vuong, TN Le, TD Ho, TNM Ung, DH Van, NH Nguyen.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Ethics approval

This study received approval from the Ethics Committee at the University of Medicine and Pharmacy in Ho Chi Minh City (approval number: 166/HDDD-ĐHYD, February 26th, 2021).

REFERENCES

1. Aluru JS, Barsouk A, Saginala K, Rawla P, Barsouk A. Valvular heart disease epidemiology. *Med Sci.* 2022;10(2):32.
2. Mensah GA, Roth GA, Fuster V. The global burden of cardiovascular diseases and risk factors: 2020 and beyond. *J Am Coll Cardiol.* 2019;74(20):2529-32.
3. Cazaubiel I, Iung B. 0433: Evolution of mitral organic valve disease in Vietnam during last two decades. *Arch Cardiovasc Dis Suppl.* 2015;7(1):50-1.
4. Otto CM, Nishimura RA, Bonow RO, Carabello BA, Erwin JP 3rd, Gentile F, et al. 2020 ACC/AHA guideline for the management of patients with valvular heart disease: a

- report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2021;143(5):e72-e227.
5. Vo AT, Nguyen DH, Van Hoang S, Le KM, Nguyen TT, Nguyen VL, et al. Learning curve in minimally invasive mitral valve surgery: a single-center experience. *J Cardiothorac Surg*. 2019;14(1):213.
 6. Anyanwu AC, Itagaki S, Chikwe J, El-Eshmawi A, Adams DH. A complexity scoring system for degenerative mitral valve repair. *J Thorac Cardiovasc Surg*. 2016;151(6):1661-70.
 7. Li J, Zhao Y, Zhou T, Zhu K, Zhai J, Sun Y, et al. Learning curve of mitral valve repair: cumulative sum failure analysis from single high-volume center. *J Thorac Dis*. 2020;12(11):6563-72.
 8. Hage F, Hage A, Manian U, Tzemos N, Chu MWA. Left ventricular remodeling after mini-mitral repair—does the complexity of mitral disease matter? *J Card Surg*. 2019;34(10):913-8.
 9. Tanaka K, Ohtaki E, Matsumura T, Misu K, Tohbaru T, Asano R, et al. Impact of a preoperative mitral regurgitation scoring system on outcome of surgical repair for mitral valve prolapse. *Am J Cardiol*. 2003;92(11):1306-9.
 10. Malhotra A, Siddiqui S, Wadhawa V, Pandya H, Patel K, Shah K, et al. ‘CLAS’ score: an objective tool to standardize and predict mitral valve reparability. *Indian J Thorac Cardiovasc Surg*. 2019;35(1):15-24.
 11. Nakayama T, Nakamura Y, Yasumoto Y, Yoshiyama D, Kuroda M, Nishijima S, et al. Early and mid-term outcomes of minimally invasive mitral valve repair via right mini-thoracotomy: 5-year experience with 129 consecutive patients. *Gen Thorac Cardiovasc Surg*. 2021;69(8):1174-84.
 12. Wan S, Lee APW, Jin CN, Wong RHL, Chan HHM, Ng CSH, et al. The choice of mitral annuloplasty ring—beyond “surgeon’s preference”. *Ann Cardiothorac Surg*. 2015;4(3):261-5.
 13. Scott Rankin J, Gaca JG, Brunsting LA 3rd, Daneshmand MA, Milano CA, Glower DD, et al. Increasing mitral valve repair rates with nonresectional techniques. *Innovations*. 2011;6(4):209-20.
 14. Marin-Cuartas M, Davierwala PM. Minimally invasive mitral valve surgery. In: Raja SG, editor. *Cardiac surgery*. Cham: Springer; 2020. p. 429-36.
 15. Perier P, Hohenberger W, Lakew F, Batz G, Urbanski P, Zacher M, et al. Toward a new paradigm for the reconstruction of posterior leaflet prolapse: midterm results of the “respect rather than resect” approach. *Ann Thorac Surg*. 2008;86(3):718-25.
 16. Castillo JG, Anyanwu AC, El-Eshmawi A, Adams DH. All anterior and bileaflet mitral valve prolapses are repairable in the modern era of reconstructive surgery. *Eur J Cardio-thorac Surg*. 2014;45(1):139-45.
 17. Koprivanac M, Kelava M, Alansari S, Javadikasgari H, Tappuni B, Mick S, et al. Degenerative mitral valve disease—contemporary surgical approaches and repair techniques. *Ann Cardiothorac Surg*. 2017;6(1):38-46.
 18. Alfieri O, Denti P. Alfieri stitch and its impact on mitral clip. *Eur J Cardio-thorac Surg*. 2011;39(6):807-8.
 19. Papadimitraki ED, Patrianakos A, Pitsis A, Marketou M, Zacharaki A, Parthenakis F. Mitral commissural prolapse. *Echocardiography*. 2021;38(4):646-56.
 20. Kawashima T, Okamoto K, Wada T, Shuto T, Umeno T, Miyamoto S. Femoral artery anatomy is a risk factor for limb ischemia in minimally invasive cardiac surgery. *Gen Thorac Cardiovasc Surg*. 2021;69(2):246-53.
 21. Lamelas J, Williams RF, Mawad M, LaPietra A. Complications associated with femoral cannulation during minimally invasive cardiac surgery. *Ann Thorac Surg*. 2017;103(6):1927-32.
 22. Bui VDA, Le MM, Nguyen D, Pham CTV, Thomas H, Nguyen DH. Compartment syndrome following minimally invasive mitral valve repair: a case report. *SAGE Open Med Case Rep*. 2022;10:2050313X221135995.
 23. David TE, David CM, Tsang W, Lafreniere-Roula M, Manlihot C. Long-term results of mitral valve repair for regurgitation due to leaflet prolapse. *J Am Coll Cardiol*. 2019;74(8):1044-53.
 24. Randolph Chitwood W Jr. Mitral repair failures are not the robot’s fault! *J Thorac Cardiovasc Surg*. 2018;155(1):e17-8.